

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 182 262 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
27.02.2002 Bulletin 2002/09

(51) Int Cl.⁷: **C12P 17/04**, C12N 1/20
// (C12N1/20, C12R1:01)

(21) Application number: 01119585.6

(22) Date of filing: 16.08.2001

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 23.08.2000 EP 00118059

(71) Applicant: **Roche Vitamins AG**
4070 Basel (CH)

(72) Inventors:

- **Asakura, Akira**
Fujisawa-shi, Kanagawa-ken 251-0032 (JP)
- **Hoshino, Tatsuo**
Kamakura-shi, Kanagawa-ken 248-0027 (JP)
- **Shinjoh, Masako**
Kamakura-shi, Kanagawa-ken 247-0061 (JP)

(74) Representative: **Müller, Ingrid, Dr. et al**
Grenzacherstrasse 124
CH-4070 Basel (CH)

(54) **Microbial production of l-ascorbic acid and d-erythorbic acid**

(57) A process for producing L-ascorbic acid or D-erythorbic acid or in each case its sodium, potassium or calcium salt, from 2-keto-L-gulonic acid or 2-keto-D-gluconic acid or in each case its sodium, potassium or calcium salt, involves incubating 2-keto-L-gulonic acid or 2-keto-D-gluconic acid, each as the free acid or as its sodium, potassium or calcium salt, and cells of a thermoacidophilic microorganism at temperatures from about 30°C to about 100°C and at a pH from about 1 to

about 6 in a solution to form L-ascorbic acid or D-erythorbic acid or an appropriate salt thereof, and isolating said product from the solution. L-Ascorbic acid (vitamin C) is widely used not only in health care as such but also in food, animal feed, e.g. fish feed, and cosmetics. D-erythorbic acid is mainly used as an antioxidant for food additives.

EP 1 182 262 A1

Description

[0001] The present invention concerns a novel microbial process for producing L-ascorbic acid and D-erythorbic acid and salts thereof. More specifically, the present invention concerns a novel process for producing L-ascorbic acid or D-erythorbic acid from 2-keto-L-gulonic acid or 2-keto-D-gluconic acid, respectively, or the respective salts from appropriate salts of the starting materials, using cells of a thermoacidophilic microorganism. L-Ascorbic acid (vitamin C) is widely used not only in health care as such but also in food, animal feed, e.g. fish feed, and cosmetics. D-erythorbic acid is mainly used as an antioxidant for food additives.

[0002] L-Ascorbic acid has been produced from D-glucose by the well-known Reichstein method (Helv. Chim. Acta 17, 311-328, 1934). In this multi-step method, L-ascorbic acid is produced chemically from the intermediate 2-keto-L-gulonic acid. The method has been used commercially for more than 60 years with many chemical and technical modifications to improve the efficiency of each of the steps to the intermediates D-sorbitol, L-sorbose, diacetone-L-sorbose, diacetone-2-keto-L-gulonic acid, 2-keto-L-gulonic acid and methyl 2-keto-L-gulonate, and to the final product L-ascorbic acid. The conversion of D-sorbitol to L-sorbose is the sole microbial step, the others being chemical steps. The conversion of diacetone-2-keto-L-gulonic acid into L-ascorbic acid has been performed by two different procedures: 1) deprotection to give 2-keto-L-gulonic acid, followed by esterification with methanol and base-catalyzed cyclization, and 2) acid-catalyzed cyclization to L-ascorbic acid directly from the protected or deprotected 2-keto-L-gulonic acid. These conversion processes must be performed in non-aqueous or low-aqueous reaction media. Environmentally and economically, reaction without organic solvents is preferable.

[0003] In its turn, D-erythorbic acid has been produced from D-glucose via 2-keto-D-gluconic acid, which itself can be produced by fermentation with a strain belonging to the genus *Pseudomonas*, and via methyl 2-keto-D-gluconate.

[0004] Much time and effort has been devoted to finding other methods of producing L-ascorbic acid using microorganisms. Most studies on the microbial production of L-ascorbic acid have focused on the production of the intermediate 2-keto-L-gulonic acid, in particular from L-sorbose (G. Z. Yin et al., Sheng Wu Hsueh Pao. 20, 246-251, 1980; A. Fujiwara et al., EP 213 591; T. Hoshino et al., US Patent No. 4,960,695; and I. Nogami et al., EP 221 707), from D-sorbitol (A. Fujiwara et al., EP 213 591; T. Hoshino et al., US Patent No. 5,312,741; M. Niwa et al., WO 95/23220; and S. F. Stoddard et al., WO 98/17819), or from D-glucose via 2,5-diketogluconic acid with a single, mixed or recombinant culture (T. Sonoyama et al., Appl. Environ. Microbiol. 43, 1064-1069, 1982; and S. Anderson et al., Science 230, 144-149, 1985). The 2-keto-L-gulonic acid can then be converted into L-ascorbic acid by chemical means as described above.

[0005] The involvement of a biological process for the conversion of 2-keto-L-gulonic acid ester into L-ascorbic acid has recently been reported in US Patent No. 6,022,719 (J.C. Hubbs). In this patent, there is described a process for producing L-ascorbic acid by contacting 2-keto-L-gulonic acid or an ester thereof with a hydrolase enzyme catalyst selected from the group consisting of a protease, an esterase, a lipase and an amidase. The patent exemplifies the formation of L-ascorbic acid from an ester of 2-keto-L-gulonic acid such as butyl 2-keto-L-gulonate, but not the formation of L-ascorbic acid from 2-keto-L-gulonic acid itself. It discloses, for example, that *Candida antarctica* B lipase catalyzed the reaction to form 413 to 530 mg/l of methyl 2-keto-L-gulonate, but no L-ascorbic acid, from 1% (w/v) 2-keto-L-gulonic acid in the presence of 8.6% methanol at a pH of from 3.1 to 3.2 and at 38°C. Ester synthetic activity of *Candida antarctica* B lipase on 2-keto-L-gulonic acid, an α -keto-carboxylic acid, at acidic pH was apparently positive, but intramolecular ester formation by this lipase was negligible.

[0006] In addition to the hydrolase reaction, ester bond syntheses such as those for proteins (amino-esters), for fatty acid esters (carboxyl-esters) and for nucleotide chains (phospho-esters) are highly functional in cells. Even in aqueous phase, the reactions proceed unidirectionally and are seldom inhibited by the product, compared with the reverse reaction of a hydrolase. Such reaction systems require a supply of activated esters such as activated transfer ribonucleic acid (tRNA), adenosine triphosphate (ATP), acyl coenzyme A (acyl-CoA) and the like, which are generated through energy-converting metabolism in cells. "In vitro" reconstitution of the reactions needs a stoichiometric supply or regeneration system of energy donors such as ATP, which are expensive for commercial production of vitamins and fine chemicals such as L-ascorbic acid and D-erythorbic acid. Commercially, utilization of intact cells is one of the preferred methods.

[0007] As mentioned above, the chemical conversion of 2-keto-L-gulonic acid to L-ascorbic acid via 2-keto-L-gulonic acid γ -lactone is an acid-catalyzed reaction accompanied by the elimination of a water molecule. The main principle of the reaction is a carboxyl ester bond formation to afford a γ -lactone ring in a 2-keto-L-gulonic acid molecule. Therefore, especially in the aqueous phase, the final state in the equilibrium reaction is determined by the physico-chemical conditions. Productivity of L-ascorbic acid from 2-keto-L-gulonic acid by chemical conversion is considerable even in aqueous phase, but not sufficient for commercial application. Performing the process in aqueous phase or in an aqueous phase with a low content of an organic solvent is highly desirable for cost effectiveness and for complying with environmental demands. Accordingly, some biological enhancement of the chemical conversion is desirable for the production in aqueous phase. Clearly, both high temperature and acidic (low) pH are desirable reaction parameters.

for improving the efficiency of the chemical reaction. But, in general, such physico-chemical conditions are known to be biologically incompatible for cell living and/or cellular activity of most of the microorganisms viable under mesophilic conditions. Utilization of thermophilic or acidophilic microorganisms is well known, but there have been few examples of the use of thermoacidophilic microorganisms which have tolerance to both heat and acid.

[0008] Surprisingly, it has now been found that the conversion of 2-keto-L-gulonic acid, as the free acid or as its sodium potassium or calcium salt, to L-ascorbic acid or the appropriate salt in aqueous phase can be directly and favorably performed by thermoacidophilic microorganisms under biologically extreme conditions, i.e. at high temperature and low (acid) pH. It has further been surprisingly found that the conversion of 2-keto-D-gluconic acid, as the free acid or as its sodium, potassium or calcium salt, to D-erythorbic acid or the appropriate salt in aqueous phase can also be directly and favorably performed by thermoacidophilic microorganisms under biologically extreme conditions.

[0009] Accordingly, the present invention provides a process for producing L-ascorbic acid or its sodium, potassium or calcium salt from 2-keto-L-gulonic acid or its sodium, potassium or calcium salt, or for producing D-erythorbic acid or its sodium, potassium or calcium salt from 2-keto-D-gluconic acid or its sodium, potassium or calcium salt, using thermoacidophilic microorganisms. More particularly, the process of the present invention is a process for producing L-ascorbic acid or its sodium, potassium or calcium salt from 2-keto-L-gulonic acid or its sodium, potassium or calcium salt, or for producing D-erythorbic acid or its sodium, potassium or calcium salt from 2-keto-D-gluconic acid or its sodium, potassium or calcium salt, comprising incubating 2-keto-L-gulonic acid or 2-keto-D-gluconic acid, each as the free acid or as its sodium, potassium or calcium salt, and cells of a thermoacidophilic microorganism which is capable of producing and/or enhancing the production of L-ascorbic acid or its sodium, potassium or calcium salt from 2-keto-L-gulonic acid or its sodium, potassium or calcium salt, or respectively D-erythorbic acid or its sodium, potassium or calcium salt, at a high temperature, i.e. at temperatures from about 30°C to about 100°C, and at an acidic pH, i.e. at a pH from about 1 to about 6, in a solution to form L-ascorbic acid or its sodium, potassium or calcium salt or D-erythorbic acid or its sodium, potassium or calcium salt, and isolating L-ascorbic acid or its sodium, potassium or calcium salt or D-erythorbic acid or its sodium, potassium or calcium salt from the solution.

[0010] A further embodiment of the present invention is a microorganism that produces L-ascorbic acid or a salt thereof or D-erythorbic acid or a salt thereof having the following characteristics:

- (a) an rDNA sequence that is at least 98.1% identical to SEQ ID NOs: 1, 2 or 3 using the Genetyx-SV/R software program;
- (b) a rod-shaped morphology;
- (c) a width of about 0.8 μm ;
- (d) an inability to grow under anaerobic conditions;
- (e) exhibiting catalase activity;
- (f) ω -Cycohexylic acid as its major fatty acid;
- (g) an ability to grow at a pH of 3.0 and a temperature of 60°C;
- (h) an inability to grow under the following conditions:

pH	Temperature
3.0	30°C
6.5	60°C
6.5	30°C

- (i) an ability to produce a (1) L-ascorbic acid or a salt thereof from 2-keto-L-gulonic acid or a salt thereof, (2) D-erythorbic acid or a salt thereof from 2-keto-D-gluconic acid or a salt thereof, or (3) both L-ascorbic acid or a salt thereof and D-erythorbic acid or a salt thereof from 2-keto-L-gulonic acid or a salt thereof and 2-keto-D-gluconic acid or a salt thereof, respectively.

[0011] For the sake of brevity "or its sodium, potassium or calcium salt" or an equivalent expression as applied to "2-keto-L-gulonic acid", "2-keto-D-gluconic acid", "L-ascorbic acid" or "D-erythorbic acid" will be mainly referred to hereinafter in each case as "or its salt". Moreover, any given concentrations of these acids or their salt forms will be expressed as being based on the free acid form even though a salt form may be present, unless clearly stated for the acid or its particular salt form which is present.

[0012] In the present description the term "thermophilic microorganism" generally means a microorganism which has optimal growth at a temperature above about 55°C, and the term "acidophilic microorganism" means a microorganism which has optimal growth at a pH in the acidic range, particularly below about pH 6, and no growth in the neutral area, i.e. in the pH range from about 6 to about 8. Thus in the context of the present invention the term "thermoacidophilic

microorganism" means a microorganism which has both these properties, i.e. which has optimal growth at a temperature above about 55°C, and at a pH below about 6 and no growth in the pH range from about 6 to about 8. The term "thermoacidophilic microorganism" also includes for the purposes of the present invention mutants of a thermoacidophilic microorganism.

5 [0013] The term "growth" as used in the present invention means that a colony formation can be observed after 20 hours of incubation. The term "no growth" as used in the present invention means that no colonies are observed after incubation for 20 hours.

[0014] Normally, thermoacidophilic microorganisms exist in and are isolable from prokaryotes and are classified under both *Archaea* and *Bacteria*. In the *Archaea* domain, the genera *Sulfolobus* (T. D. Brock et al., Arch. Mikrobiol. 84, 54-68, 1972) and *Thermoplasma* (M. DeRosa et al., Phytochemistry 170, 1416-1418, 1970) are well-known as thermoacidophilic microorganisms, and the genera *Acidanus* (A. H. Segerer et al., Int. J. Syst. Bacteriol. 36, 559-564, 1986), *Desulfurolobus* (W. Zilling et al., Syst. Appl. Microbiol. 8, 197-209, 1986), *Metallosphaera* (G. Huber et al., Syst. Appl. Microbiol. 12, 38-47, 1989), *Picrophilus* (C. Schleper et al., J. Bacteriol. 177, 7050-7059, 1995) and *Stygiolobus* (A. H. Segerer et al., Int. J. Syst. Bacteriol. 41, 495-501, 1991) have also been reported as being thermoacidophilic microorganisms of the *Archaea* domain. In the *Bacteria* domain, the genera *Acidimicrobium* (D. A. Clark et al., Microbiology 142, 785-790, 1996), *Acidotherrmus* (F. Rainey et al., FEMS Microbiol. Lett., 108, 27-30, 1993), *Sullobacillus* (R. S. Golovacheva et al., Microbiology 47, 658-665, 1978) and *Alicyclobacillus* (G. Darland et al., J. Gen. Microbiol. 67, 9-15, 1971; G. Deinhard et al., Syst. Appl. Microbiol. 10, 47-53, 1987) are thermoacidophilic microorganisms

[0015] The thermoacidophilic microorganisms useable in the present invention are any thermoacidophilic microorganisms which are capable of producing and/or enhancing the production of L-ascorbic acid or its salt from 2-keto-L-gulonic acid or its salt, or D-erythorbic acid or its salt from 2-keto-D-gluconic acid or its salt.

[0016] The thermoacidophilic microorganisms used in the present invention can be obtained from any kind of natural sources thereof, including soils and hot spring water, and from artificial sources thereof, including processed acidic foods and beverages, e.g. fruit juices and mixed fruit/vegetable juices.

25 [0017] The more extreme the conditions, i.e. the higher the temperature and the lower (more acid) the pH, under which any particular thermoacidophilic microorganism displays tolerance, the more preferably this microorganism is used in the process of the present invention. Besides tolerance to heat and acidity, thermoacidophilic microorganisms which are also tolerant to a high concentration, i.e. from about 5% to about 20% (w/v), of 2-keto-L-gulonic acid or its salt, or of 2-keto-D-gluconic acid or its salt, in solution when incubated at high temperature and acidic pH are also more preferably used in the process. In addition, thermoacidophilic microorganisms with the aerobic and chemoorganotrophic characteristics described in the present invention are preferable for the efficient, i.e. rapid, production of cells.

[0018] Preferred thermoacidophilic microorganisms are those derived from prokaryotes, including bacteria and archaea. More preferred microorganisms are thermoacidophilic bacteria. Especially preferred thermoacidophilic microorganisms are bacteria belonging to the genus *Alicyclobacillus*.

35 [0019] Among the thermoacidophilic bacteria, the genus *Alicyclobacillus* embraces most of the strictly aerobic, spore-forming, rod-shaped and chemoorganotrophic bacteria. These microorganisms were initially assigned to the genus *Bacillus*. However, phylogenetic analysis based on sequence comparisons of the 16S rRNA gene has shown that the genus *Alicyclobacillus* belongs to a distinct line of descent within the low G + C Gram-positive lineage of the *Bacillus* (J. D. Wisotzkey et al., Int. J. Syst. Microbiol. 42, 263-269, 1992). The three validly named species of the genus *Alicyclobacillus* (A.) are: *A. acidocaldarius* (DSM 446^T, G. Darland et al., J. Gen. Microbiol. 67, 9-15, 1971), *A. acidoterrestis* (DSM 3922^T, G. Deinhard et al., Syst. Appl. Microbiol. 10, 47-53, 1987) and *A. cycloheptanicus* (DSM 4006^T, G. Deinhard et al., Syst. Appl. Microbiol. 10, 68-73, 1987). Besides sequence comparisons of the 16S rRNA genes, the most distinguishable characteristic of these microorganisms is the possession of structural units of ω -cyclohexyl fatty acids (ω -cyclohexylundecanoic acid, ω -cyclohexyltridecanoic acid) or of ω -cycloheptyl fatty acids (ω -cycloheptylundecanoic acid, ω -cycloheptyltridecanoic acid) (L. Albuquerque et al., Int. J. Syst. Evol. Microbiol. 50, 451-457, 2000). Several strains with the characteristics of the genus *Alicyclobacillus* have been isolated so far from acidic soils within geothermal areas and from certain non-geothermal soils. In addition to soil samples, they have also been isolated from many acidic beverages as spoilage bacteria (G. Cerny et al., Z Lebens Unters Forsch 179, 224-227, 1984; K. Yamazaki et al., Biosci. Biotech. Biochem. 60, 543-545, 1996; M. Niwa et al., Japanese Patent Publication (Kokai) No. 140696/1996). Recently, a wide diversity of genospecies among the genus *Alicyclobacillus* has been proposed in addition to the three validly named species (A. Hiraishi et al., J. Gen. Appl. Microbiol. 43, 295-304, 1997; L. Albuquerque et al., Int. J. Syst. Evol. Microbiol. 50, 451-457, 2000)

[0020] Preferred thermoacidophilic microorganisms used in the present invention have the following characteristics:

55 1) Thermoacidophilic growth:

Showing growth at pH 3.0 at 60°C in 20 hours, but showing no growth at pH 3.0 at 30°C, at pH 6.5 at 30°C and at pH 6.5 at 60°C, in each case in 20 hours.

2) ω -cyclohexyl fatty acids.

Possessing ω -cyclohexyl fatty acid structural units according to gas chromatography-mass spectrometry (GC/MS) analysis.

3) 16S rRNA sequence similarity

Phylogenic analysis of 16S genes coding for rRNA sequences confirms the allocation to the genus *Alicyclobacillus*.

[0021] The thermoacidophilic microorganisms used in the present invention can be obtained from natural and artificial sources, as indicated above, or commercially from culture depositories. For isolating the microorganisms from such natural and artificial sources the appropriate microorganism source, such as a natural source soil or hot spring water, or an artificial source processed acidic food or beverage, is conveniently cultured in an aqueous medium and/or on a solid medium supplemented with appropriate nutrients under aerobic conditions. The cultivation is conveniently conducted at temperatures above about 40°C and at pHs below about 5, preferably above about 50°C and below about pH 4, and more preferably above about 55°C and below about pH 3.5. While the cultivation period varies depending upon the pH, temperature and nutrient medium used, a period of 12 hours to several days will generally give favorable results.

[0022] Especially preferred thermoacidophilic microorganisms belonging to the genus *Alicyclobacillus* and used in the present invention are *Alicyclobacillus* sp. DSM No. 13652 and DSM No. 13653 which are obtainable from the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Mascheroder Weg 1b, D-38124 Braunschweig, Germany, and the new strains, *Alicyclobacillus* sp. NA-20 and NA-21, which were isolated from a soil sample collected at Iwate Prefecture, Japan, and *Alicyclobacillus* sp. FJ-21, which was isolated from a commercial acidic beverage, in particular a fruit juice, purchased at Kamakura-shi, Kanagawa Prefecture, Japan.

[0023] All five especially preferred thermoacidophilic microorganisms were deposited under the Budapest Treaty on August 17, 2000 at the DSMZ and were allotted the following accession numbers:

Alicyclobacillus sp. : DSM No. 13652
Alicyclobacillus sp. : DSM No. 13653
Alicyclobacillus sp. NA-20 : DSM No. 13649
Alicyclobacillus sp. NA-21 : DSM No. 13650
Alicyclobacillus sp. FJ-21 : DSM No. 13651

[0024] The last three mentioned thermoacidophilic microorganisms, being new, represent a further aspect of the present invention.

[0025] In the present invention, and as indicated above, mutants of the above mentioned thermoacidophilic microorganisms can also be used. The mutant of a microorganism used in the present invention can be induced by treating a wild type strain with a mutagen such as irradiation by ultraviolet rays, X-rays, γ -rays or contact with nitrous acid or other suitable mutagens. A mutant may also be obtained by isolating a clone occurring by spontaneous mutation thereof, which may be effected by methods known by skilled artisans for such purposes. Many of these methods have been described in specialized publications, for example "Chemical Mutagens" edited by Y. Tajima, T. Yoshida and T. Kada, Kodansha Scientific Inc., Tokyo, Japan, 1973.

[0026] Moreover, biologically and taxonomically homogeneous cultures of *Alicyclobacillus* sp. DSM No. 13652, DSM No. 13653, NA-20 (DSM No. 13649), NA-21 (DSM No. 13650) or FJ-21 (DSM No. 13651) can be used. Said term "biologically and taxonomically homogeneous cultures" in the present invention means cultures showing the following biological and taxonomical characteristics:

- growth: aerobic and thermoacidophilic
- spore: forming
- cell morphology: rod-shaped
- major fatty acids: ω -cyclohexyl fatty acids
- phylogenetical position: most close (more than 90% identity in the nucleotide sequence of 16S rRNA gene) to the strains classified in the genus *Alicyclobacillus*, such as *Alicyclobacillus* sp. DSM No. 13652, *A. sp.* UZ-1, *A. sp.* MIH-2, *A. sp.* KHA-31 and *A. acidocaldarius* DSM446^T, whereby the identity in the nucleotide sequence is defined using the Nucleotide Sequence Homology program (Genetyx-SV/R, version 4.0, Software Development Co., Tokyo, Japan) with default conditions (unit size to compare = 1)

[0027] The above mentioned thermoacidophilic microorganisms can be used in any form, in particular as intact cells, modified cells or immobilized cells. Methods for immobilizing cells are well known in the art (see for example W. M. Fogarty et al., *Microbial Enzymes and Biotechnology*, 2nd Edition, Elsevier Applied Science, pp. 373-394 (1983), and Japanese Patent Publication No. 61265/1994).

[0028] The screening of thermoacidophilic microorganisms to assess their suitability for use in the process of the present invention can be performed by the following method:

[0029] The appropriate microorganism source is cultured in an aqueous medium containing the substrate 2-keto-L-gulonic acid or its salt or 2-keto-D-gluconic acid or its salt and supplemented with appropriate nutrients under moderately aerobic conditions, i.e. under aerobic incubation without enforced aeration or vigorous agitation. A concentration of the substrate, 2-keto-L-gulonic acid or its salt, or 2-keto-D-gluconic acid or its salt, for carrying out the cultivation may be from about 3% (w/v) to about 20% (w/v), preferably from about 4% (w/v) to about 18% (w/v), and more preferably from about 5% (w/v) to about 16% (w/v). The incubation may be conducted at pHs from about 0.5 to about 4.0, preferably from about 1.0 to about 3.5, and more preferably from about 1.5 to about 3.0, and at temperatures from about 45°C to about 90°C, preferably from about 50°C to about 85°C, and more preferably from about 55°C to about 80°C. While the incubation period varies depending upon the pH, temperature and nutrient medium used, a period of about 12 hours to several days will generally give favorable results. After the incubation, the suitability of the use of the screened thermoacidophilic microorganism in the process of the present invention may be assessed by its degree of productivity (high accumulation of the product) of L-ascorbic acid or its salt, or D-erythorbic acid or its salt, compared with the productivity in a "blank" incubation, i.e. without the thermoacidophilic microorganism, whereas an over 2-fold increase of productivity over the productivity in a "blank" incubation is preferred.

[0030] In the incubation described above, the presence of a high concentration of the substrate, 2-keto-L-gulonic acid or its salt, or 2-keto-D-gluconic acid or its salt, in addition to a high temperature and an acidic pH, may represent an extreme physico-chemical condition even for a thermoacidophilic microorganism. The tolerance to 2-keto-L-gulonic acid or its salt, or 2-keto-D-gluconic acid or its salt, at high temperature and an acidic pH may be an important characteristic of the thermoacidophilic microorganisms used in the process of the present invention for retention (maintenance) of the living cell state.

[0031] The incubation for the production and/or enhancing the production of L-ascorbic acid or its salt from 2-keto-L-gulonic acid or its salt, or D-erythorbic acid or its salt from 2-keto-D-gluconic acid or its salt, with cells of a thermoacidophilic microorganism in the process of the present invention is effected in solution in an aqueous phase. The solvent for the aqueous phase is preferably water alone, i.e. without any added other solvent(s). If an additional solvent is used, however, this is preferably a lower alkanol such as methanol. The incubation for the production and/or enhancing the production of L-ascorbic acid from 2-keto-L-gulonic acid, or D-erythorbic acid from 2-keto-D-gluconic acid, each of these products or substrates being present as the free acid or the respective sodium, potassium or calcium salt, requires nutrients such as assimilable carbon sources, digestible or assimilable nitrogen sources and inorganic substances, trace elements, vitamins, L-amino acids and other growth promoting factors. As assimilable carbon sources, D-glucose, sucrose, D-glucono- δ -lactone, starch and the like can be employed. Various organic or inorganic substances may be employed as nitrogen sources, such as yeast extract, meat extract, peptone, casein, corn steep liquor, urea, amino acids, nitrates, ammonium salts such as ammonium sulfate, and the like. As inorganic substances magnesium sulfate, potassium phosphate, sodium chloride, potassium chloride, calcium chloride, and the like may be employed. Furthermore, as trace elements sulfates, hydrochlorides or phosphates of calcium, magnesium, zinc, manganese, cobalt and iron may be employed. Particularly suitable as inorganic salts are monopotassium phosphate, magnesium sulfate, ferrous sulfate and manganese sulfate. And, if necessary, conventional nutrient factors or an antifoaming agent, such as animal oil, vegetable oil or mineral oil, can be added.

[0032] The conditions of the incubation may vary depending on the species and generic character of the thermoacidophilic microorganism employed. The incubation is effected at what is considered to be a high temperature for an incubation, i.e. at temperatures from about 30°C to about 100°C, preferably from about 40°C to about 95°C, and most preferably from about 55°C to about 95°C, at an acidic pH, particularly at a pH from about 1.0 to about 6.0, preferably from about 1.0 to about 4.5, and most preferably from about 1.5 to about 3.0, under an aerobic condition. Normally, an incubation period ranging from about 1 to about 100 hours is sufficient.

[0033] The suitable initial concentration of 2-keto-L-gulonic acid or its salt or of 2-keto-D-gluconic acid or its salt, as appropriate, for the incubation depends on the particular thermoacidophilic microorganism used. However, a concentration of 2-keto-L-gulonic acid or its salt, or of 2-keto-D-gluconic acid or its salt, in each case based on the (equivalent) amount of free acid, of from about 5% (w/v) to about 20% (w/v) is generally used. This concentration is preferably from about 10% (w/v) to about 15% (w/v).

[0034] The process of the present invention shows the following characteristics:

a) Specific production rate of L-ascorbic acid or its salt:

The specific production rate for L-ascorbic acid is, for example (in the presence of 8% (w/v) 2-keto-L-gulonic acid, the substrate, and 2.5 g/L L-ascorbic acid, the product, at 59°C and pH 2.5 for 20 hours by the strain NA-21 (DSM No. 13650)) estimated to be about 2.3 mg of L-ascorbic acid / mg of crude cellular proteins / hour. This is based on the results given in Example 7, hereinafter.

b) Product inhibition.

The production in the present invention is seldom inhibited by the product, L-ascorbic acid or its salt, or D-erythorbic acid or its salt. The production in the process of the present invention may provide higher conversion yield than those obtained by reversible reactions in aqueous phase.

[0035] The L-ascorbic acid or its salt, or D-erythorbic acid or its salt, formed in solution may be separated and purified by conventional methods known in the art. If the product is the sodium, potassium or calcium salt of the respective acid, this salt may, if desired, be converted into the respective free acid by conventional methods known in the art. In each case isolation of the product may be effected by methods relying upon the differences in properties between the product and impurities (including the non-converted substrate) such as solubility, adsorbability, an electrochemical property and the distribution coefficient between two solvents. Use of an absorbent such as an ion exchange resin is an example of a convenient process for the isolation of the product. Use of an electro-dialysis system is a further example of a convenient process for isolation of the product. If the product is insufficiently pure for its subsequent use, it may be purified by conventional methods such as recrystallization and chromatography.

[0036] L-Ascorbic acid can be produced from L-sorbose or D-sorbitol by combination of the organism having the activity for converting 2-keto-L-gulonic acid to L-ascorbic acid or 2-keto-D-gluconic acid to D-erythorbic acid with an organism having L-sorbose/L-sorbose dehydrogenase and D-sorbitol dehydrogenase and which can convert D-sorbitol and/or L-sorbose to 2-keto-L-gulonic acid (see A. Fujiwara et al., EP 213 591; T. Hoshino et al., US Patent No. 4,960,695; T. Hoshino et al., US Patent No. 5,312,741); ex. *Gluconobacter oxydans* DSM 4025 in one-step conversion with one vessel or two-step conversion with two vessels.

[0037] D-Erythorbic acid can be produced from D-glucose or D-gluconic acid by combination of the organism having the activity for converting 2-keto-L-gulonic acid to L-ascorbic acid or 2-keto-D-gluconic acid to D-erythorbic acid with an organism having D-glucose dehydrogenase (Ameyama et al., Agric Biol. Chem. 45:851-861, 1981) and/or D-gluconate dehydrogenase (Shinagawa et al., Agric Biol. Chem. 48: 1517-1522, 1984); ex. *Gluconobacter dioxyacetonicus* IFO 3271 and which can convert D-glucose and/or D-gluconic acid to 2-keto-D-gluconic acid in one-step conversion with one vessel or two step-conversion with two vessels.

[0038] The following Examples are illustrate the process of the present invention and are not intended to limit the scope of the invention in any way.

Example 1

Screening of thermoacidophilic microorganisms:

A) Isolation from soil samples

[0039] Soil samples collected at an acidic hot spring area at Iwate Prefecture in Japan were used for the screening. Thermoacidophilic microorganisms were recovered from the soil samples in 0.9% (w/v) NaCl solution, and isolated by spreading the solution on 573c (pH 3.5) agar plate medium containing 0.1% (w/v) of D-glucose, 0.13% (w/v) of $(\text{NH}_4)_2\text{SO}_4$, 0.1% (w/v) of yeast extract, 0.15% (w/v) of KH_2PO_4 , 0.025% (w/v) of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.007% (w/v) of $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ and 2% (w/v) of agar (the pH was adjusted by 6N H_2SO_4 ; D-glucose and agar were sterilized separately). After incubation at 60°C for 20 hours, single colonies grown at pH 3.5 and 60°C were randomly collected and purified by transferring them three times onto the same agar plate medium under the same conditions. The resulting isolated microorganisms were numbered and designated as belonging to the "NA" series. For example, one such isolated microorganism was designated as NA-20, another as NA-21.

B) Isolation from acidic beverages

[0040] Various commercial acidic beverages, i.e. fruit juice products and mixed fruit/vegetable juice products, were subjected to experiments for isolating thermoacidophilic microorganisms therefrom. In each case 1 ml of the commercial product was centrifuged and the resulting pellet was washed with 1 ml of sterilized distilled water to obtain a washed pellet. The pellet was suspended in 0.1 ml of sterilized distilled water, and the suspension was spread on an agar plate of medium 573c. The plate was then incubated at 60°C for 1 to 3 days to observe single colonies, which were purified by transferring them three times onto the same agar plate medium under the same conditions. The resulting isolated microorganisms were numbered and designated as belonging to the "FJ" series. For example, one such isolated microorganism was designated as FJ-21.

Example 2

Screening of NA- and FJ-series isolated microorganisms for production of L-ascorbic acid from 2-keto-L-gulonic acid

- 5 [0041] Screening for production of L-ascorbic acid from 2-keto-L-gulonic acid was performed by the following living cell reaction system. Compositions of LM101 type media, mineral mixture [MM], vitamin mixture [VM] and amino acid mixture [AM] are listed in Tables 1 below.

Tables 1

Table 1a: Media (salt base)

	LM101c	LM101d
D-glucose	variable	variable
(NH ₄) ₂ SO ₄	4	1.3
KH ₂ PO ₄	1.5	1.5
MgSO ₄ ·7H ₂ O	0.25	0.25
NaCl	0.1	0.1
KCl	0.1	0.1
CaCl ₂ ·2H ₂ O	0.07	0.07
[g/L]		
pH	2.5 ^a	2.5 ^b

a: after addition of sodium 2-keto-L-gulonate monohydrate, pH is adjusted by 6N H₂SO₄ after addition of 2-keto-L-gulonic acid, pH is adjusted by 6N KOH

Table 1b

Mineral Mixture [x 1000 conc.]	
CuSO ₄ ·5H ₂ O	620
FeSO ₄ ·5H ₂ O	600
MnSO ₄ ·5H ₂ O	600
ZnSO ₄ ·5H ₂ O	600
CoCl ₂ ·5H ₂ O	790
Na ₂ MoO ₄ ·5H ₂ O	700
[mg/L]	

Table 1c

Vitamin Mixture [x 100 conc.]	
Biotin	100
Ca(+)-Pantothenate	100
Folic acid	100
Inositol	200
Pyridoxal phosphate·H ₂ O	100
Riboflavin	10

Table 1c (continued)

Vitamin Mixture [x 100 conc.]	
Thiamine-HCl	100
Nicotinamide	100
[mg/L]	

Table 1d

L-Amino Acid Mixture [x 5 conc.]	
Arginine-HCl	630
Cystine	120
Glutamine	1460
Histidine-HCl-H ₂ O	210
Isoleucine	260
Leucine	260
Lysine-HCl	360
Methionine	76
Phenylalanine	165
Threonine	240
Tryptophan	50
Tyrosine	180
Valine	230
[mg/L]	

[0042] The thermoacidophilic microorganisms of the NA and FJ series, isolated as described in Example 1, were grown on 573c agar plate medium (pH 3.5, 60°C, 15 hours) and inoculated into medium LM101c-plus ["plus" means supplementation with MM (x 1 conc.), VM (x 0.1 conc.) and AM (x 0.01 conc.)] containing 0.25% (w/v) of D-glucose. After aerobic cultivation in test tubes (60°C, 8 hours), the resulting cells were collected by centrifugation, and were used as seeding cells after suspending in LM101c salt base (about 25 of optical density at 660 nm [OD₆₆₀]). The cells were inoculated at the final OD₆₆₀ of 0.2 into 0.8 ml of LM101c-plus medium (pH 2.5) containing 6% (w/v) of sodium 2-keto-L-gulonate monohydrate and 0.2% (w/v) of D-glucose. The resulting cell reaction mixture was incubated at 59°C under a moderately aerobic condition; the test tube (2.0 ml micro test tube, Eppendorf, Germany) was incubated with a pin hole (0.65 mm) at the top with rotary shaking (120 rpm with 45 mm radius). L-Ascorbic acid production was determined by HPLC analysis: YMC-Pack Polyamine II column (ID. 4.6 x 150 mm; YMC Co., Japan) at 264 nm with the mobile-phase solvent containing 70% (v/v) of acetonitrile and 15 mM of ammonium dihydrogen phosphate at a flow rate of 1.5 ml/min. The amount of water lost by evaporation was estimated as the weight decrease during the incubation, and a compensing amount of water was added to retain the original volume. After 23 hours cultivation, the L-ascorbic acid production of each strain was compared with two control values: "aerobic blank" being the L-ascorbic acid production obtained with the same medium and conditions but without cell inoculation, and "anaerobic control" being the L-ascorbic acid production with the same medium but without cell inoculation in a closed micro test tube with argon gas. As a result of the screening, strains NA-20, NA-21 and FJ-21 were selected as potent L-ascorbic acid producers; they produced 1.07, 1.19 and 1.23 g/L of L-ascorbic acid, respectively. The amounts of L-ascorbic acid in the aerobic blank and anaerobic control were about 0.10 and 0.30 g/L, respectively.

[0043] The living-cell reaction with 2-keto-D-gluconic acid (hemicalcium salt containing 1.5 mol/mol H₂O, SIGMA Chemical Co., St. Louis, MO, USA) was performed under the same conditions as described above except using 1.2% (w/v) of 2-keto-D-gluconic acid instead of 6% (w/v) of sodium 2-keto-L-gulonate monohydrate as the substrate. The production of D-erythorbic acid was determined by the same HPLC analysis method. The strain NA-21 produced 0.051 g/L of D-erythorbic acid after 23 hours incubation. The amounts of D-erythorbic acid in the aerobic blank and anaerobic control were not detectable (less than 0.001 g/L).

Example 3

Taxonomy of the isolated microorganisms

- 5 [0044] The three isolated microorganisms, NA-20, NA-21 and FJ-21, were aerobic, spore-forming and rod-shaped bacteria. Phenotypic characters of the isolates are summarized in the following Table 2.

Table 2

	NA-20	NA-21	FJ-21
10 Shape	rod	rod	rod
Size			
(width, μm)	0.8	0.8	0.8
(length, μm)	2-3	3-5	2-3
15 Gram Stain	variable	variable	negative
Motility	+	+	-
Anaerobic Growth	-	-	-
20 Oxidase Test	-	-	-
Catalase Test	+	+	+
Major Fatty Acid	ω -Cyclohexylic acids (C17 and 19)	ω -Cyclohexylic acids (C17 and 19)	ω -Cyclohexylic acids (C17 and 19)
25 Growth at ^a			
pH Temp. ($^{\circ}\text{C}$)			
3.0 60	+	+	+
3.0 30	-	-	-
6.5 60	-	-	-
30 6.5 30	-	-	-

a: growth on 573c agar plate for 20 hours

- 35 [0045] The microorganisms showed growth at pH 3.0/60 $^{\circ}\text{C}$ in 20 hours, but not at pH 3.0/30 $^{\circ}\text{C}$, pH 6.5/30 $^{\circ}\text{C}$ and pH 6.5/60 $^{\circ}\text{C}$ in 20 hours. Therefore, they were characterized as thermoacidophilic *Bacillus* group bacteria. GC/MS analysis on fatty acids indicated that the major components of the three isolated microorganisms were identical with those of *Alicyclobacillus acidocaldarius* DSM 466^T investigated as the control, suggesting that ω -cyclohexyl fatty acids were the major components of the isolated microorganisms as well as of *Alicyclobacillus acidocaldarius* DSM 466^T. 16S rRNA gene sequences of the isolated microorganisms were determined with 16S rRNA gene kit (PE Applied Biosystems, USA; SEQ ID NOs: 1, 2 and 3 for the strain NA-20, NA-21 and FJ-21, respectively) and were subjected to the BLAST search program (J. Mol. Biol. 215 403-410, 1990; Nucleic Acids Res. 25 3389-3402, 1997). The sequence similarity analysis using the Nucleotide Sequence Homology Program (Genetyx-SV/R, version 4.0, Software Development Co., Tokyo, Japan) with default conditions (unit size to compare = 1) indicated that the isolated microorganisms could belong phylogenetically to the genus *Alicyclobacillus*. The following Table 3 shows the levels of binary sequence identity (%) in 16SrRNA gene sequences between the isolated microorganisms and reference strains including three type strains.

Table 3

Identity (%) in 16SrRNA gene						
		Accession No	NA-20	NA-21	FJ-21	
50	<i>A. acidocaldarius</i>	DSM 446 ^T	X60742	96.9	97.0	98.3
	<i>A. acidoterrestris</i>	DSM 3922 ^T	AJ133631	94.0	94.1	94.5
	<i>A. cycloheptanicus</i>	DSM 4006 ^T	X51928	92.6	92.6	92.9
55	<i>A. sp.</i>	UZ-1	AB004579	99.5	99.6	97.8
	<i>A. sp.</i>	MIH-2	AB004580	99.5	99.6	97.8
	<i>A. sp.</i>	KHA-31	AB004581	99.2	99.3	97.6

Table 3 (continued)

Identity (%) in 16SrRNA gene					
		Accession No.	NA-20	NA-21	FJ-21
A. sp.	DSM 13652	AJ133634	99.0	99.1	97.6
A. sp.	NA-20		100.0	99.7	98.1
A. sp.	NA-21		99.7	100.0	98.1
A. sp.	FJ-21		98.1	98.1	100.0

[0046] The 16S rDNA sequences of NA-20 and NA-21 were most similar to each other (99.7%) and to the sequences of *Alicyclobacillus* sp. UZ-1, MIH-2, KHA-31 (J. Gen. Appl. Microbiol., 43, 295-304, 1997) and *A. sp.* DSM 13652 (99.0 to 99.6%). The sequence of FJ-21 was most similar to those of *A. acidocaldarius* DSM 446^T, NA-20 and NA-21 (98.1 to 98.3%). From these results, the three isolated microorganisms were classified into the genus *Alicyclobacillus* and named as *Alicyclobacillus* sp. NA-20, NA-21 and FJ-21, respectively.

[0047] Under the same screening conditions as described in Example 2, *Alicyclobacillus* sp. DSM 13652 and 13653 produced 0.91 and 0.95 g/L of L-ascorbic acid from sodium 2-keto-L-gulonate monohydrate, respectively.

Example 4

Effects of carbon/energy source and aeration

[0048] In the L-ascorbic acid production process described in Example 2, the medium containing 0.2% (w/v) of D-glucose in addition to 6% (w/v) of sodium 2-keto-L-gulonate monohydrate was used to maintain cell living. For the production using *Alicyclobacillus* sp. NA-21, addition of D-glucono- δ -lactone [0.1% (w/v)], sucrose [0.1% (w/v)] or soluble starch [1% (w/v)] instead of D-glucose resulted in almost the same production of L-ascorbic acid, but no addition of carbon/energy source resulted in no additional production over the aerobic blank (see Table 4).

Table 4

Carbon	[% (w/v)]	Cell	L-Ascorbic Acid [g/L] ^a
Sucrose	0.1	+	0.69
Soluble Starch	1.0	+	0.65
D-Glucono- δ -lactone	0.1	+	0.76
D-Glucose	0.1	+	0.80
D-Glucose	0.025	+	0.66
No Carbon	-	+	0.10
Aerobic Blank	-	-	0.09

a: 13 hour production from 6% (w/v) of sodium 2-keto-L-gulonate monohydrate

[0049] In the L-ascorbic acid production process described in Example 2, the production was carried out under moderately aerobic condition for cell living. Using *Alicyclobacillus* sp. NA-21, the results of carrying out the process under aerobic and anaerobic conditions were compared. The same reaction mixture in a completely closed micro test tube was used for the system under anaerobic conditions after gasification with argon gas. The aerobic conditions contributed to nearly linear production until 38 hours, but anaerobic conditions did not allow production even after 15 hours (see Tables 5).

Tables 5

L-Ascorbic acid production (g/L) under aerobic condition				
	Reaction Time [h]			
	0	15	23	38
+ Cell	0.00	0.80	1.29	2.10
- Cell	0.00	0.12	0.13	0.17

L-Ascorbic acid production (g/L) under anaerobic condition				
	Reaction Time [h]			
	0	15	23	38
+ Cell	0.00	0.79	0.87	1.05
- Cell	0.00	0.25	0.36	0.59

Example 5

L-Ascorbic acid production by using the strain NA-21 or its derivatives

[0050] L-Ascorbic acid production from 6% (w/v) of sodium 2-keto-L-gulonate monohydrate was examined by using *Alicyclobacillus* sp. NA-21. Preparation of seeding cell was carried out by the same method as described in Example 2. The cells were inoculated at the final OD₆₆₀ of 0.25 into 0.8 ml of LM101c-plus medium (pH 2.5) containing 6% (w/v) of sodium 2-keto-L-gulonate monohydrate and 0.1% (w/v) of D-glucose. The resulting cell reaction mixture was incubated at 59°C under a moderately aerobic condition; the test tube was incubated with a pin hole (0.65 mm) at the top with rotary shaking (120 rpm with 45 mm radius). L-Ascorbic acid production by *Alicyclobacillus* sp. NA-21 continued linearly until 38 hours to reach to 2.23 g/L of L-ascorbic acid. Productivity was obviously more effective than that of the aerobic blank and of the anaerobic control (see Tables 6).

Tables 6

OD 660				
	Reaction Time [h]			
	0	15	23	38
+ Cell	0.25	0.22	0.20	0.15
- Cell Aerobic Blank	0.00	0.00	0.00	0.00
Anaerobic Control	0.00	0.00	0.00	0.00

pH				
	Reaction Time [h]			
	0	15	23	38
+ Cell	2.46	2.46	2.45	n.d.
- Cell Aerobic Blank	2.46	2.44	2.43	n.d.
Anaerobic Control	2.46	2.44	2.44	n.d.
n.d.: not determined				

L-Ascorbic Acid Production [g/L]				
	Reaction Time [h]			
	0	15	23	38
+ Cell	0.00	0.76	1.25	2.23
- Cell Aerobic Blank	0.00	0.12	0.13	0.17
Anaerobic Control	0.00	0.25	0.36	0.59

[0051] Two derivatives from the original strain, *Alicyclobacillus* sp. NA-21, MA-10 and MB-6 were sequentially obtained by a conventional mutation followed by the selection steps described below. From the original strain, MA-10 was selected as a strain having tolerance against higher 2-keto-L-gulonic acid concentration (apparently about 1% (w/v) improvement from 10% (w/v) at 60°C and pH 2.5) after ultraviolet irradiation. Subsequently, from MA-10, MB-6 was

selected as a strain having tolerance against higher temperature (apparently about 2°C improvement from 60°C at 11% (w/v) of 2-keto-L-gulonic acid and pH 2.5) after treatment with N-methyl-N'-nitro-N-nitrosoguanidine. L-Ascorbic acid productivities were compared among the original strain, MA-10 and MB-6 by the same living-cell reaction described above at 62°C and pH 2.5, using 11% (w/v) of 2-keto-L-gulonic acid and medium containing x 0.5 conc. of LM101d salt base, 0.25% (w/v) of D-glucose, x 0.1 conc. of MM, x 0.05 conc. of VM and x 0.005 conc. of AM. The original strain, MA-10 and MB-6 produced 2.02, 2.22 and 2.80 g/L of L-ascorbic acid at 23 hours, respectively. The aerobic blank and the anaerobic control produced 1.63 and 1.94 g/L of L-ascorbic acid, respectively.

Example 6

L-Ascorbic acid production from 8% (w/v) 2-keto-L-gulonic acid with feeding of cells and D-glucose

[0052] L-Ascorbic acid production from 8% (w/v) 2-keto-L-gulonic acid with feeding of cells and D-glucose was examined by using *Alicyclobacillus* sp. NA-21. Preparation of seeding cell was carried out by the same method as described in Example 2. The cells were inoculated at the final OD660 of 0.25 into 0.8 ml of LM101d-plus medium (pH 2.5) containing 8% (w/v) of 2-keto-gulonic acid and 0.15% (w/v) of D-glucose. The resulting cell reaction mixture was incubated at 59°C under a moderately aerobic condition; the test tube was incubated with a pin hole (0.65 mm) at the top with rotary shaking (120 rpm with 45 mm radius). Seeding cells and D-glucose were supplied at 24 hours (0.25 of OD660 and 0.15% (w/v) at final concentration, respectively) as a spot feeding. L-Ascorbic acid production by *Alicyclobacillus* sp. NA-21 continued linearly until 47 hours to reach 3.70 g/L of L-ascorbic acid (see Table 7).

Table 7

L-Ascorbic Acid Production [g/L]								
Reaction Time (h)	0	15	23	39	47	71	87	95
+ Cell	0.00	1.13	1.97	2.99	3.70	n.d.	n.d.	n.d.
- Cell Aerobic Blank	0.00	0.23	0.32	0.35	0.36	0.39	0.32	0.28
Anaerobic Control	0.00	0.49	0.74	1.20	1.52	2.26	2.57	2.44
n.d.: not determined								

[0053] On the other hand, productivity of the anaerobic control reached a maximum at about 2.5 g/L of L-ascorbic acid after about 80 hours.

Example 7

L-Ascorbic acid production from 8% (w/v) 2-keto-L-gulonic acid in the presence of L-ascorbic acid.

[0054] L-Ascorbic acid production from 8% (w/v) 2-keto-L-gulonic acid in the presence of 0 - 4.5 g/L of L-ascorbic acid was examined by using *Alicyclobacillus* sp. NA-21. Preparation of seeding cell was carried out by the same method as described in Example 2. The cells were inoculated at the final OD660 of 0.25 into 0.8 ml of LM101d-plus medium (pH 2.5) containing x 0.1 conc. of MM instead of x 1 conc. of MM and 8% (w/v) of 2-keto-gulonic acid and 0.20% (w/v) of D-glucose. The resulting cell reaction mixture was incubated at 59°C under moderately aerobic conditions; the test tube was incubated with a pin hole (0.65 mm) at the top with rotary shaking (120 rpm with 45 mm radius) for 20 hours. To the above medium, 0.0, 1.1, 2.3 or 4.5 g/L of L-ascorbic acid (product) was added at 0 hour (before the production). Productivity of *Alicyclobacillus* sp. NA-21 was nearly the same in the absence and presence of the product. On the other hand, productivity of the anaerobic control was gradually repressed by the presence of a higher concentration of the product (see Tables 8).

Tables 8

From 0.0 g/L of Initial L-Ascorbic Acid		
	L-Ascorbic Acid [g/L] Reaction Time [h]	Net Production * 20 h minus 0 h
	0	20
		[g/L]
+ Cell	0	1.81
- Cell Aerobic Blank	0	0.44

EP 1 182 262 A1

Tables 8 (continued)

From 0.0 g/L of Initial L-Ascorbic Acid		
	L-Ascorbic Acid [g/L] Reaction Time [h]	Net Production * 20 h minus 0 h
	0 20	[g/L]
Anaerobic Control	0 0.83	0.83

From 1.3 g/L of Initial L-Ascorbic Acid		
	L-Ascorbic Acid [g/L] Reaction Time [h]	Net Production 20 h minus 0 h
	0 20	[g/L]
+ Cell	1.32 3.16	1.84
- Cell Aerobic Blank	1.32 0.80	-0.52
Anaerobic Control	1.32 1.75	0.43

From 2.5 g/L of Initial L-Ascorbic Acid		
	L-Ascorbic Acid [g/L] Reaction Time [h]	Net Production 20 h minus 0 h
	0 20	[g/L]
+ Cell	2.48 4.33	1.85
- Cell Aerobic Blank	2.48 1.55	-0.93
Anaerobic Control	2.48 2.7	0.22

From 4.6 g/L of Initial L-Ascorbic Acid		
	L-Ascorbic Acid [g/L] Reaction Time [h]	Net Production 20 h minus 0 h
	0 20	[g/L]
+ Cell	4.55 6.64	2.09
- Cell Aerobic Blank	4.55 3.02	-1.53
Anaerobic Control	4.55 4.21	-0.34
*: Net Production means amount of L-ascorbic acid accumulated at 20 h minus that added at 0 h.		

Sequence Listing

5 <110> Roche Vitamins AG
 <120> Microbial process for producing L-ascorbic acid and D-erythorbic acid
 <130> Alicyclobacillus NA20, 21, FJ21 16S nuc
 10 <140>
 <141>
 <150> EP Application No. 00118059.5
 <151> 2000-08-23
 15 <160> 3
 <170> PatentIn Ver. 2.1
 <210> 1
 <211> 1529
 20 <212> DNA
 <213> Alicyclobacillus sp.
 <220>
 <221> rRNA
 <222> (1)..(1529)
 25 <223> NA-20 partial 16SrRNA gene sequence
 <400> 1
 agagtttgat cctggctcag gacgaacgct ggcggcgtgc ctaatacatg caagtcgagc 60
 ggggtctctc ggaggccagc ggcggacggg tgaggaacac gtgggtaatc tgccctttcag 120
 gccggaataa cgcccgaaa cgcccgctaa tgccggatac gcccgcgagg aggcattctc 180
 30 ttgcggggga aggcccaatt ggcgcgctga gagaggagcc cgccgcat tagctngttg 240
 gcggggtaac ggcacacaa ggcgacgat cgtagccgac ctgagagggt gaccggccac 300
 actgggactg agacacggcc cagactccta cgggaggcag cagtaggga tcttccgcaa 360
 tggcgcaag cctgacggag caacgcgcg tgagcgaaga aggccttcg gttgtaaagc 420
 tctgttctc ggggagagcg gcatggggga tggaaagccc catgcgagac ggtaccgagt 480
 gaggaagccc cggctaacta cgtgccagca gccgcggtaa aacgtagggg gcgagcgttg 540
 35 tccggaatca ctgggcgtaa aggggtgcgt ggcggtcgag caagtctgga gtgaaagtcc 600
 atggctcaac catgggatgg ctttggaac tgcctgactt gagtgtgga gaggcaagg 660
 gaattccacg tgtagcgtg aaatgcgtag agatgtggag gaataccagt ggcgaaggcg 720
 ccttgctgga cagtgactga cgctgagcga cgaaagcgtg gggagcaaac aggattagat 780
 accctggtag tcacgcgcgt aaacgatgag tgctaggtgt tggggggaca caccctcagt 840
 ccgaaggaaa cccaataagc actccgcctg gggagtacgg tcgcaagact gaaactcaa 900
 40 ggaattgacg ggggcccgca caagcagtgg agcatgtggt ttaattcgaa gcaaccgcaa 960
 gaaccttacc agggcttgac atccctctga cacnctcaga gatgaggggg cccttcgggg 1020
 cagaggagac aggtggtgca tggttgtcgt cagctcgtgt cgtgagatgt tgggttcagt 1080
 ccgcaacga gcgcaacct tgacctgtgt taccagcgcg ttgaggcggg gactcacagg 1140
 tgactgcccg cgtaatcgcg aggaaggcgg ggatgacgtc aaatcatcat gccctgatg 1200
 tcctgggcta cacacgtgct acaatggcg gaacaaaggg agcgaagcc gcgaggcgga 1260
 45 gcgaaaccca aaaagccgct cgtagttcgg attgcaggc gcaactcggc tgcataaagc 1320
 cggaattgct agtaatcgcg gatcagcat ccgcggtgaa tacgttccc ggccttgta 1380
 acaccgcccg tcacaccacg agagtcggca acaccgaag tcggtgaggt aaccctnng 1440
 gggagccagc cgccgaaggt ggcgtcgatg attggggtga agtcgtaaca aggtagcgt 1500
 accggaaggt gcggtcgat cactcctt 1529
 50 <210> 2
 <211> 1529
 <212> DNA
 <213> Alicyclobacillus sp.
 55 <220>
 <221> rRNA

<222> (1)..(1529)

<223> NA-21 Partial 16SrRNA gene

<400> 2

```

5 agagtttcat cctggctcag gacgaacgct ggccggctgc ctaatacatg caagtcgagc 60
  gggctctcttc ggaggccagc ggccggacggg tgaggaaacac gtgggtaatc tgcctttcag 120
  gccggaataa cgcccggaac cgccggctaa tgccggatac gcccgcgagg aggcattctc 180
  ttgcggggga aggcccaatt ggccactga gagaggagcc cgcggcgcat tagctngttg 240
  gcggggtaac ggcccaccaa ggccagcatg cgtagccgac ctgagagggt gaccggccac 300
  actgggactg agacacggcc cagactccta cgggaggcag cagtagggaa tcttcgcgaa 360
10 tggcgcaag cctgacggag caacgccgcg tgagcgaaga aggccttcgg gttgtaaaagc 420
  tctgttgctc ggggagagcg gcatggggga tggaaaagccc catgcgagac ggtaccgagt 480
  gaggaagccc cggctaacta cgtgccagca gcccggttaa aacgtagggg gcgagcgttg 540
  tccggaatca ctggcgctaa aggtgtcgta ggccgtcgag caagtctgga gtgaaagtc 600
  atggtcaca catgggatgg ctttgaaaac tgcttgactt gaggctgga gaggcaagg 660
  gaattccacg tgtagcgggt aaatcgctag agatgtggag gaataccagt ggcgaaggcg 720
15 ccttgctgga cagtactga cgtgaggca cgaagcgtg gggagcaaac aggattagat 780
  accctggtag tccacgccgt aaacgatgag tgctaggtgt tggggggaca caccacagtg 840
  ccgaaggaaa cccaataagc actccgcctg gggagacgg tcgcaagact gaaactcaa 900
  ggaattgacg ggggcccgca caagcagtg agcatgtgtt ttaattcgaa gcaacgcgaa 960
  gaaccttacc agggcttgac atccctctga caccctcaga gatgaggggg cccttcgggg 1020
  cagaggagac aggtgttgca tgggtgtcgt cagctcgtgt cgtgagatgt tgggttcagt 1080
20 cccgcaacga gcgcaaccct tgacctgtgt taccagcgcg ttgaggcggg gactcacagg 1140
  tgactgccg tgtaagtccg aggaaggcgg ggtgacgtc aaatcatcat gccctgatg 1200
  tcttgggcta cacacgtgct acaatggcg gaacaaagg aggcgaagcc gcgaggcgga 1260
  gcgaaaccca aaaagccgct cgtagtccg attgcaggct gcaactcgc tgcatgaagc 1320
  cggaattgct agtaatcgcg gatcagcatg ccgcgtgtaa tacgttccg ggccttgatc 1380
  acaccgcccg tcacaccacg agagtcgcga acaccgaaag tcggtgaggt aaccccttag 1440
25 gggagccagc cgccgaagggt cgggtcgatg attgggtgta agtcgtaaca aggtagccgt 1500
  accggaaggt gcggctggat cacctcctt 1529

```

<210> 3

<211> 1495

30 <212> DNA

<213> Alicyclobacillus sp.

<220>

<221> rRNA

<222> (1)..(1495)

35 <223> FJ-21 Partial 16SrRNA gene sequence

<400> 3

```

  aggacgaacg ctggcggcgt gcctaataca tgcaagtcga gggacacctt tctgaggtca 60
  gcggcggagc ggtgaggaac acgtgggtaa tctgccttcc agaccggaat aacgcccgga 120
  aacggggcgt aatgccggat acgcccgcga ggaggcatct tcttgcgggg aaaggcccga 180
  ttgggcccgt gagagaggag cccgcggcgc attagctngt tggcggggta acggcccacc 240
  aaggcgacga tgcgtagccg acctgagagg gtgaccggcc acactgggac tgagacacgg 300
  cccagactcc tacgggaggg agcagtaggg aatcttccgc aatgggcgca agcctgacgg 360
  agcaacgcgg cgtgagcgaa gaaggccttc ggggtgtaaa gctctgttgc tcggggagag 420
  cgccatgggg agtggaagc cccatgcgag acggtaccga gtgagggaag cccggctaac 480
  tacgtgccag cagccgcggg aaaacgtagg gggcgagcgt tgcgggaat cactgggcgt 540
45 aaagggtgcg taggcggtcg agcaagtcgt gaggtaaggt ccatgggtca accatgggat 600
  ggctctggaa actgcttgac ttgagtgctg gagaggcaag ggggaattcca cgtgtagcgg 660
  tgaaatgcgt agagatgtgg aggaatacca gtggcgagg cgccttgctg gacagtgaat 720
  gacgctgagg cacgaaagcg tggggagcaa acaggattag ataccctggt agtccacgcc 780
  gtaaacgatg agtgctaggt gttgggggga cacaccccag tgccgaagga aacccaataa 840
  gcactccgcc tggggagtag ggtcgcaaga ctgaaactca aagggaattga cggggggccc 900
50 cacaagcagt ggagcatgtg gtttaattcg aagcaacgcg aagaacctta ccagggcttg 960
  acatccctct gacgggtgca gagatgcacc ttcccttcgg gcgagaggag acaggtgggt 1020
  catggttgct gtcagctcgt gtcgtgagat gttgggttca gtcccgaac gagcgcaacc 1080
  cttgacctgt gttaccagcg cgnatangcg gggactcaca ggtgactgcc ggcgtaagtc 1140
  ggagggaagg ggggatgacg tcaaatcatc atgccctga tgcctgggc tacacacgtg 1200
  ctacaatggg cggtacaaag ggaggcgaa cgcgaggcgg gagcgaaacc caaaaagccc 1260
55 ctcgtagtgc ggattgcagg ctgcaactcc cctgcatgaa gccggaattg ctagttaatc 1320
  cggatcagca tgcccggtg aatacgttcc cgggccttgt acacaccgcc cgtcacacca 1380

```


cgagagtcgg caacacccga agtcggtgag gtaaccccga aaggggagcc agccgccgaa 1440
 ggtggggtcg atgattgggg tgaagtcgta acaaggtagc cgtaccggaa ggtgc 1495

Claims

1. A process for producing L-ascorbic acid or its sodium, potassium or calcium salt from 2-keto-L-gulonic acid or its sodium, potassium or calcium salt, or D-erythorbic acid or its sodium, potassium or calcium salt from 2-keto-D-gluconic acid or its sodium, potassium or calcium salt, comprising incubating 2-keto-L-gulonic acid or 2-keto-D-gluconic acid, each as the free acid or as its sodium, potassium or calcium salt, and cells of a thermoacidophilic microorganism at temperatures from about 30°C to about 100°C and at a pH from about 1 to about 6 in a solution to form L-ascorbic acid or D-erythorbic acid or an appropriate salt thereof, and isolating said L-ascorbic acid, D-erythorbic acid or appropriate salt thereof from the solution.
2. The process according to claim 1 wherein the thermoacidophilic microorganism is isolated from prokaryotes.
3. The process according to claim 2 wherein the prokaryotes from which the thermoacidophilic microorganism is isolated are classified under bacteria.
4. The process according to claim 3 wherein the thermoacidophilic microorganism belongs to the bacteria genus *Alicyclobacillus*.
5. The process according to claim 4 wherein the thermoacidophilic microorganism of the genus *Alicyclobacillus* is *Alicyclobacillus* sp. DSM No. 13652, DSM No. 13653, NA-20 (DSM No. 13649), NA-21 (DSM No. 13650) or FJ-21 (DSM No. 13651) or in each case a mutant thereof.
6. The process according to claim 4 wherein the thermoacidophilic microorganism of the genus *Alicyclobacillus* is a biologically and taxonomically homogeneous culture having the identifying characteristics of *Alicyclobacillus* sp. DSM No. 13652, DSM No. 13653, NA-20 (DSM No. 13649), NA-21 (DSM No. 13650) or FJ-21 (DSM No. 13651).
7. The process according to any one of claims 1 to 6 wherein the solution contains water as the solvent.
8. The process according to any one of claims 1 to 7 wherein the incubation is effected under an aerobic condition.
9. The process according to any one of claims 1 to 8 wherein the incubation is effected under an aerobic condition and in the presence of nutrients.
10. A process according to any one of claims 1 to 9 wherein the concentration of the substrate, 2-keto-L-gulonic acid, 2-keto-D-gluconic acid or in each case the sodium, potassium or calcium salt thereof in the solution is from about 5% (w/v) to about 20% (w/v), preferably from about 10% (w/v) to about 15% (w/v), based on the amount of free acid.
11. A process according to any one of claims 1 to 10 wherein the incubation is effected at a temperature from about 40 to about 95°C, preferably from about 55 to about 95°C.
12. A process according to any one of claims 1 to 11 wherein the incubation is effected at a pH from about 1.0 to about 4.5, preferably from about 1.5 to about 3.0.
13. *Alicyclobacillus* sp. NA-20 (DSM No. 13649), *Alicyclobacillus* sp. NA-21 (DSM No. 13650) and *Alicyclobacillus* sp. FJ-21 (DSM No. 13651).



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 01 11 9585

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	EP 1 026 257 A (HOFFMANN LA ROCHE) 9 August 2000 (2000-08-09) * the whole document *	1-3,7-12	C12P17/04 C12N1/20 //(C12N1/20, C12R1:01)
D,A	WO 97 43433 A (EASTMAN CHEM CO) 20 November 1997 (1997-11-20) * examples 14,15 *	1	
D,X	WISOTZKEY J D ET AL: "COMPARATIVE SEQUENCE ANALYSES ON THE 16S RRNA RDNA OF BACILLUS-ACIDOCALDARIUS BACILLUS-ACIDOTERRESTRIS AND BACILLUS-CYCLOHEPTANICUS AND PROPOSAL FOR CREATION OF A NEW GENUS ALICYCLOBACILLUS NEW-GENUS" INTERNATIONAL JOURNAL OF SYSTEMATIC BACTERIOLOGY, vol. 42, no. 2, 1992, pages 263-269, XP001041904 ISSN: 0020-7713 * the whole document *	13	
D,X	HIRAISHI AKIRA ET AL: "Phylogenetic characterization of a new thermoacidophilic bacterium isolated from hot springs in Japan." JOURNAL OF GENERAL AND APPLIED MICROBIOLOGY, vol. 43, no. 5, October 1997 (1997-10), pages 295-304, XP001041901 ISSN: 0022-1260 * the whole document *	13	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 12 December 2001	Examiner Lejeune, R
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 82 (P04001)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 11 9585

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

12-12-2001

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
EP 1026257	A	09-08-2000	BR	0000073 A	26-09-2000
			CN	1263950 A	23-08-2000
			EP	1026257 A1	09-08-2000
			JP	2000210094 A	02-08-2000
			US	6146860 A	14-11-2000
WO 9743433	A	20-11-1997	US	5817490 A	06-10-1998
			AU	3075697 A	05-12-1997
			BR	9709099 A	03-08-1999
			CN	1225686 A	11-08-1999
			EP	0938582 A1	01-09-1999
			JP	2001505042 T	17-04-2001
			WO	9743433 A1	20-11-1997
			US	6022719 A	08-02-2000
			US	6136575 A	24-10-2000
			US	6271006 B1	07-08-2001

EPO FORM P449

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

cgagagtcgg caacacccga agtcggtgag gtaaccccgaa aaggggagcc agccgccgaa 1440
 ggtggggtcg atgattgggg tgaagtcgta acaaggtagc cgtaccggaa ggtgc 1495

5

10

Claims

1. A process for producing L-ascorbic acid or its sodium, potassium or calcium salt from 2-keto-L-gulonic acid or its sodium, potassium or calcium salt, or D-erythorbic acid or its sodium, potassium or calcium salt from 2-keto-D-gluconic acid or its sodium, potassium or calcium salt, comprising incubating 2-keto-L-gulonic acid or 2-keto-D-gluconic acid, each as the free acid or as its sodium, potassium or calcium salt, and cells of a thermoacidophilic microorganism at temperatures from about 30°C to about 100°C and at a pH from about 1 to about 6 in a solution to form L-ascorbic acid or D-erythorbic acid or an appropriate salt thereof, and isolating said L-ascorbic acid, D-erythorbic acid or appropriate salt thereof from the solution.
2. The process according to claim 1 wherein the thermoacidophilic microorganism is isolated from prokaryotes.
3. The process according to claim 2 wherein the prokaryotes from which the thermoacidophilic microorganism is isolated are classified under bacteria.
4. The process according to claim 3 wherein the thermoacidophilic microorganism belongs to the bacteria genus *Alicyclobacillus*.
5. The process according to claim 4 wherein the thermoacidophilic microorganism of the genus *Alicyclobacillus* is *Alicyclobacillus* sp. DSM No. 13652, DSM No. 13653, NA-20 (DSM No. 13649), NA-21 (DSM No. 13650) or FJ-21 (DSM No. 13651) or in each case a mutant thereof.
6. The process according to claim 4 wherein the thermoacidophilic microorganism of the genus *Alicyclobacillus* is a biologically and taxonomically homogeneous culture having the identifying characteristics of *Alicyclobacillus* sp. DSM No. 13652, DSM No. 13653, NA-20 (DSM No. 13649), NA-21 (DSM No. 13650) or FJ-21 (DSM No. 13651).
7. The process according to any one of claims 1 to 6 wherein the solution contains water as the solvent.
8. The process according to any one of claims 1 to 7 wherein the incubation is effected under an aerobic condition.
9. The process according to any one of claims 1 to 8 wherein the incubation is effected under an aerobic condition and in the presence of nutrients.
10. A process according to any one of claims 1 to 9 wherein the concentration of the substrate, 2-keto-L-gulonic acid, 2-keto-D-gluconic acid or in each case the sodium, potassium or calcium salt thereof in the solution is from about 5% (w/v) to about 20% (w/v), preferably from about 10% (w/v) to about 15% (w/v), based on the amount of free acid.
11. A process according to any one of claims 1 to 10 wherein the incubation is effected at a temperature from about 40 to about 95°C, preferably from about 55 to about 95°C.
12. A process according to any one of claims 1 to 11 wherein the incubation is effected at a pH from about 1.0 to about 4.5, preferably from about 1.5 to about 3.0.
13. *Alicyclobacillus* sp. NA-20 (DSM No. 13649), *Alicyclobacillus* sp. NA-21 (DSM No. 13650) and *Alicyclobacillus* sp. FJ-21 (DSM No. 13651).



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 01 11 9585

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	EP 1 026 257 A (HOFFMANN LA ROCHE) 9 August 2000 (2000-08-09) * the whole document *	1-3,7-12	C12P17/04 C12N1/20 /(C12N1/20, C12R1:01)
D,A	WO 97 43433 A (EASTMAN CHEM CO) 20 November 1997 (1997-11-20) * examples 14,15 *	1	
D,X	WISOTZKEY J D ET AL: "COMPARATIVE SEQUENCE ANALYSES ON THE 16S RRNA RDNA OF BACILLUS-ACIDOCALDARIUS BACILLUS-ACIDOTERRESTRIS AND BACILLUS-CYCLOHEPTANICUS AND PROPOSAL FOR CREATION OF A NEW GENUS ALICYCLOBACILLUS NEW-GENUS" INTERNATIONAL JOURNAL OF SYSTEMATIC BACTERIOLOGY, vol. 42, no. 2, 1992, pages 263-269, XP001041904 ISSN: 0020-7713 * the whole document *	13	
D,X	HIRAISHI AKIRA ET AL: "Phylogenetic characterization of a new thermoacidophilic bacterium isolated from hot springs in Japan." JOURNAL OF GENERAL AND APPLIED MICROBIOLOGY, vol. 43, no. 5, October 1997 (1997-10), pages 295-304, XP001041901 ISSN: 0022-1260 * the whole document *	13	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 12 December 2001	Examiner Lejeune, R
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1501 (12/01) 28.03.2001 WPO/01

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 11 9585

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information

12-12-2001

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1026257	A	09-08-2000	BR 0000073 A	26-09-2000
			CN 1263950 A	23-08-2000
			EP 1026257 A1	09-08-2000
			JP 2000210094 A	02-08-2000
			US 6146860 A	14-11-2000
WO 9743433	A	20-11-1997	US 5817490 A	06-10-1998
			AU 3075697 A	05-12-1997
			BR 9709099 A	03-08-1999
			CN 1225686 A	11-08-1999
			EP 0938582 A1	01-09-1999
			JP 2001505042 T	17-04-2001
			WO 9743433 A1	20-11-1997
			US 6022719 A	08-02-2000
			US 6136575 A	24-10-2000
			US 6271006 B1	07-08-2001

EPO FORM P459

For more details about this annex : see Official Journal of the European Patent Office, No 12/82